

TUSSIN CF COUGH AND COLD- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution
Shopko Stores Operating Co., LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Shopko Tussin CF Cough & Cold Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

| age | dose |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 10 mL every 4 hours |
| children under 12 years | do not use |

Other information

- **each 10 mL contains:** sodium 5 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare To Robitussin® Multi-Symptom Cold active ingredients

FOR AGES 12 & OVER

ADULT

TUSSIN CF

COUGH & COLD

MULTI-SYMPTOM

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

Relieves:

Cough

Nasal Congestion

Mucus

Peak Cold

Non-Drowsy

Alcohol Free

CHERRY FLAVOR

8 FL OZ (237 mL)



TUSSIN CF COUGH AND COLD

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:370 12-516 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|----------------------------------|--------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 10 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 10 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 10 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |

Product Characteristics

| | | | |
|----------|--------|--------------|--|
| Color | RED | Score | |
| Shape | | Size | |
| Flavor | CHERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:37012-516-34 | 1 in 1 CARTON | 07/08/2013 | |
| 1 | | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 07/08/2013 | |

Labeler - Shopko Stores Operating Co., LLC (023252638)